## GLASGOW COMA SCALE : Do it this way



## Institute of Neurological Sciences NHS Greater Glasgow and Clyde



CHECK

اي تي تاب وج ڪل ڪك لماوع نا يروك اي روه لخم ريم عني د باوج - دوه يگل ريڻوچ روا



02310061

،داوم اك تيج تاب ،انلهك اك روهكنآ انرك تكرح اك مسج ريئاب روا ريئاد



ريرك مدماشم

-تساوغرد رك الي اي رك لوب :زاوآ

ندرگ ،نخان كك يلكنا ؤابد :ينامسج ـرپ يڏه يك وربااي ـمهـُپ ـمك



يدنب مجرد

قباطع عك لوعدر رتهب عس بس ـريرك ضيوفت

#### Eye opening

Criterion	Observed	Rating	Score
Open before stimulus	<b>✓</b>	Spontaneous	4
After spoken or shouted request	<b>✓</b>	To sound	3
After finger tip stimulus	<b>✓</b>	To pressure	2
No opening at any time, no interfering factor	<b>*</b>	None	1
Closed by local factor	<b>✓</b>	Non testable	NT

#### Verbal response

Criterion	Observed	Rating	Score
Correctly gives name, place and date	<b>*</b>	Orientated	5
Not orientated but communication coherently	<b>*</b>	Confused	4
Intelligible single words	<b>*</b>	Words	3
Only moans / groans	<b>*</b>	Sounds	2
No audible response, no interfering factor	<b>4</b>	None	1
Factor interferring with communication	<b>*</b>	Non testable	NT

#### Best motor response


Criterion	Observed	Rating	Score
Obey 2-part request	<b>*</b>	Obeys commands	6
Brings hand above clavicle to stimulus on head neck	<b>*</b>	Localising	5
Bends arm at elbow rapidly but features not predominantly abnormal	<b>*</b>	Normal flexion	4
Bends arm at elbow, features clearly predominantly abnormal	<b>*</b>	Abnormal flexion	3
Extends arm at elbow	<b>*</b>	Extension	2
No movement in arms / legs, no interfering factor	<b>*</b>	None	1
Paralysed or other limiting factor	<b>*</b>	Non testable	NT

## Sites For Physical Stimulation

# Supraorbital notch Finger tip pressure Trapezius Pinch

## Features of Flexion Responses

Modified with permission from Van Der Naalt 2004 Ned Tijdschr Geneeskd



**Normal flexion** 

Rapid Variable Arm away from body